



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display Abstract

Show 20

Sort by

Send to

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Protein Expr Purif. 2005 Jul 19; [Epub ahead of print]

Related Articles, Links

ELSEVIER
FULL-TEXT ARTICLE

Baculovirus expression, purification, and characterization of human protein phosphatase 2A catalytic subunits alpha and beta.

Ikehara T, Shinjo F, Ikehara S, Imamura S, Yasumoto T.

Japan Science and Technology Agency (JST) Collaboration of Regional Entities for the Advancement of Technological Excellence in Okinawa, Okinawa Health Biotechnology Research Development Center, 12-75 Suzaki, Uruma, Okinawa 904-2234, Japan.

Protein phosphatase 2A (PP2A) contains a 36-kDa catalytic subunit (PP2Ac), a 65-kDa structural subunit (PR65/A), and a regulatory B subunit. The core enzyme consists of the structural and catalytic subunits. The catalytic subunit exists as two closely related isoforms, alpha and beta. Several natural toxins, including okadaic acid (OA) and microcystins, specifically inhibit PP2A. To obtain biologically active recombinant PP2A and to compare the properties of the PP2A catalytic subunit alpha and beta isoforms, we expressed human PP2Aalpha and cbeta in High Five insect cells. The recombinant PP2Aalpha and cbeta possess similar phosphatase activities using p-NPP and phosphopeptide as substrates and are strongly inhibited by OA and microcystin-LR to similar degrees. In addition, PP2Aalpha or cbeta was co-expressed with PR65/A and co-purified as a core dimer, PP2A(D) (Aalpha/calpha and Aalpha/cbeta) with PR65alpha/Aalpha. The recombinant PP2A(D) bound to the B subunit in vitro. These results show that the recombinant PP2Aalpha and cbeta are identical in their ability to associate with the A and B subunits, in their phosphatase activities, and in carboxyl-methylation. Furthermore, our results show that High Five insect cells can produce biologically active recombinant PP2A, which should be a valuable tool for detecting natural toxins and investigating the mechanism of PP2A catalysis and other protein interactions.

PMID: 16039140 [PubMed - as supplied by publisher]